

# Salmon Escapement to Englishman River, 2006

Ian Matthews and Alan Eden<sup>1</sup>

South Coast Area  
Fisheries and Oceans Canada  
3225 Stephenson Point Road  
Nanaimo, BC  
V9T 1K3

July 2007

<sup>1</sup> – 365 Dunsmuir Rd, Qualicum Bay, V9K 2A1

## **Abstract**

From 1999 to 2003 extensive spawning salmon assessments were conducted on the Englishman River. Funding for this escapement work was provided through the Pacific Salmon Endowment Fund (PSEF). Funding and in-kind donations from The Nature Trust and Fisheries and Oceans Canada provided resources for the 2005 and 2006 escapement work. Spawning surveys are an important part of any salmon stock assessment and rebuilding program.

Bi-weekly snorkel surveys of the Englishman River mainstem were attempted through Fall 2006, however, high water flows in November and December limited the number of surveys conducted. The escapement estimate for pink salmon was 50, chinook salmon was 590, coho salmon was 490, and chum salmon was 130.

There were sources of uncertainty in data collection due to incomplete coverage, extended periods of high water when surveys were not possible and poor visibility in the lower reaches of the river. Bank erosion during high run-off was the primary source of this increased turbidity and during these high flow events it was not feasible to swim some sections of the river.

These escapements are at or above the long term average but must be considered with several factors. The methodology of escapement enumeration was changed in 1999 which would have affected the reported numbers. The marine conditions have decreased the ocean survival of smolt to returning adults, which limited the ability of the stocks to withstand high levels of exploitation by the various fisheries. As a result DFO fishery managers drastically reduced the opportunities for the commercial and sport fishing sectors for coho and chinook. This reduction in exploitation rate increased the escapement of salmon to southern BC creeks in general, not just the Englishman River.

## Introduction

Extensive salmon escapement programs were conducted from 1999 through 2003 on the Englishman River (DFO NuSEDs database, Baillie and Young 2002, Baillie and Young 2003). This work was funded through The Pacific Salmon Endowment Fund Society and the Englishman River Watershed Recovery Plan (ERWRP) steering committee. The goal of this fund is to achieve healthy, sustainable and naturally diverse salmon stocks by conserving and rebuilding salmon populations through strategic and focused efforts. The Georgia Basin coho (*Oncorhynchus kisutch*) and steelhead (*O. mykiss*) stocks are one of three concerns that the Society had initially identified as a priority for developing a recovery plan.

Due to funding constraints no escapement surveys were conducted in 2004. Funding and in-kind donations for the 2005 and 2006 escapement work were received from the Nature Trust and Fisheries and Oceans Canada.

One of the primary components of a recovery strategy is a comprehensive monitoring program. This program is used to track the salmon populations to measure whether objectives are being met and to detect stock declines and increases in each area of concern. Part of this monitoring program is to enumerate the salmon escapement using scientifically accepted practices approved by DFO.

The Englishman River watershed has all species of salmon including steelhead and is designated a sensitive stream by the BC government under the Fish Protection Act (Bocking and Gaboury 2001). Annual escapement estimates of salmon from 1953 to 2005 are presented in Table 1.

The Englishman River flows into the Strait of Georgia at Parksville on Vancouver Island and drains roughly 324 km<sup>2</sup> (Figure 1). The river originates on the eastern slopes of Mt. Arrowsmith (1820 m) and Mt. Moriarty Ridge and flows in an easterly direction for 40 km. The mainstem has an accessible reach of 15.85 km. There are four main tributaries: South Englishman River (4.5 km accessible reach), Morison Creek (2.1 km), Centre Creek (5.2 km), and Shelly Creek (1.0 km). Centre Creek is a tributary of the South Englishman, located approximately 200 m upstream from the confluence of the South Englishman with the mainstem (Bocking and Gaboury 2001).

All five species of Pacific salmon, coho, pink (*O. gorbuscha*), chum (*O. keta*), chinook (*O. tshawytscha*) and sockeye (*O. nerka*) occur in the Englishman River. As well as steelhead trout, there are rainbow trout in the system (the non-anadromous form of steelhead trout) and coastal cutthroat trout (*O. clarki clarki*). Coastrange sculpin (*Cottus aleuticus*) and prickly sculpin (*C. asper*) are also resident fish species. Other species that may be present are threespine stickleback (*Gasterosteus aculeatus*) and lamprey (*Lampetra* sp.).

This report presents the results of salmon escapement enumeration work that was done in the Englishman system in the fall of 2006.

## Methods

### Surveys

Snorkel surveys were conducted by two swimmers at approximately bi-weekly intervals beginning in early September and ending in mid January 2007 (Table 2). Surveys were confined to the Englishman River mainstem, which for the previous escapement projects had been divided into 26 sections of approximately 600 meters each. To facilitate data collection in 2006 these were combined into three survey reaches (Figure 2). The lower reach (sections 0 to 8) started at the estuary and covered the first 3.9 km to the top of Allbrook Canyon. The middle reach (sections 9 to 15) was 5.8 km long ending approximately 2 km upstream of the South Englishman confluence. The upper reach (sections 15 to 26) comprised the remaining 6.2 km to the Englishman River Falls. Two days were required for the survey crew to cover the full length of the river accessible to migrating salmon. These surveys were then combined to provide a single estimate of salmon in the river for that time period. The number of live fish observed were recorded by species.

### Population Estimate calculations

Each survey count must be expanded to account for salmon missed within the survey reach (Observer Efficiency, or OE) and for un-counted salmon outside of the survey reach (Coverage).

OE is an estimate of how many fish were missed by the swimmers and is expressed as a proportion of the total that was enumerated. This is estimated by taking into account reduced visibility from turbid water conditions, deeper pools, overcast days and amount of cover such as log jams and cut banks in which fish could hide.

Coverage refers to the proportion of the total fish population that was present in the reach surveyed by the swimmers. As the 2006 surveys were only performed on the Englishman River mainstem and were not always complete an expansion factor was applied to many of the counts to derive an estimate for the total salmon in the accessible watershed. This expansion varies by species over time and was based on survey data from the more extensive 2001 Englishman River project (Baillie and Young 2002). These data are presented in Table 3. Applying these two expansions to the observed counts from each set of surveys produced an estimate for the total number of salmon in the system at that time.

Final escapement estimates were calculated from the expanded survey data using Area-Under-the-Curve analysis (AUC) (English et al. 1992). The estimated number of fish from each survey is plotted against time and the integral of this curve is divided by the survey life (SL) of the fish. This is described by the following formulae;

1. 
$$P = \frac{\text{Count}}{\text{Obs.Eff.} \times \text{Coverage}}$$
2. 
$$AUC = \sum_{i=2}^n ((t_i - t_{i-1})(P_i + P_{i-1}) * 0.5)$$
3. 
$$\text{Esc.Est.} = \frac{AUC}{\text{SurveyLife}}$$

were  $P_i$  is the estimate of fish in the stream on the  $i^{\text{th}}$  day,  $t_i$  the number of days measured from the first day fish entered the survey area and  $n$  is the number of surveys conducted. Survey life is defined as the average length of time a salmon is available to be surveyed. When a creek is surveyed from the anadromous barrier down to the estuary, the SL used would be equal to the stream life, or the average length of time a salmon is in fresh water. If a tributary or a segment of the river were surveyed, then the SL life used would only be the average length of time a salmon is within the surveyed reach.

The survey life values derived in the more extensive 2002 Englishman River Salmon Escapement project were used for analysis of this year's data. The population estimate that resulted from the AUC calculation was rounded to two significant digits.

## Results

### ***Pink***

The estimated total escapement for pinks was 50. This number is based on the expanded peak live count that was obtained on 8<sup>th</sup> September. Observed and expanded numbers are presented in Table 3.

Pinks were only observed on the first two surveys (8<sup>th</sup> and 29<sup>th</sup> September). The escapement estimate is derived from the first survey which only covered the lower reach of the river, however, this is where the majority of the pinks were observed spawning in 2005.

### ***Chinook***

The estimated total escapement for chinook was 590. This number was derived from AUC calculations using the expanded estimates and a survey life of 18 days (Baillie and Young 2003). The peak count of 197, expanded to 276, occurred on 17<sup>th</sup> October. Observed and expanded numbers are presented in Table 3.

No chinook were observed during the first survey on 8<sup>th</sup> September and only 12 were counted on 2<sup>nd</sup> November. Continuous high water flows prevented any further surveys until 13<sup>th</sup> January. No salmonids were observed during this final survey. Chinook occurred throughout the survey area although primarily within the middle and lower reaches.

### ***Chum***

The estimated total escapement for chum was 130. This was derived from AUC calculations using the expanded estimates and a survey life of 10 days (Baillie and Young 2003). The peak count was 44 on 17<sup>th</sup> October. This was expanded to an estimated peak of 52. Observed and expanded numbers are presented in Table 3.

Chum were first observed in the river on 29<sup>th</sup> September when 5 were counted in the lower reach. Two chum was counted on 2<sup>nd</sup> November just prior to start of the Fall storms and the extended period of high water.

### ***Sockeye***

No sockeye were observed during the 2006 escapement surveys.

### ***Coho***

The estimated total escapement for coho was 490. This was derived from AUC calculations using the expanded estimates and a survey life of 20 days (Baillie and Young 2003). The peak count of 281 occurred on the 17<sup>th</sup> October survey. This was expanded to 423 as only the middle and lower reaches were covered during this survey. Observed and expanded numbers are presented in Table 3.

Coho were first observed on 29<sup>th</sup> September when 56 were counted in the lower reach. Six coho were counted on the 2<sup>nd</sup> November survey. They were observed throughout the survey area.

## Discussion

Escapement enumeration for four of the five species of Pacific salmon (pink, chum, coho and chinook) was of limited success in 2006. Above average rainfall and the frequency of winter storms moving through the region resulted in an extended period (early November to mid-January, 2007) during which high water made escapement surveys impossible (Figure 3). Consequently the escapement estimates for all four species of salmon returning to the Englishman River in 2006 are lower in accuracy and precision than those derived in 2005.

Erosion of silt from the clay banks at kilometre 8.4 continues to be a problem in the lower river. At low water flows during September and October a layer of sediment was evident for at least 500 metres downstream of the clay banks and any rain events or increase in water flows resulted in an immediate drop in visibility throughout the lower river. Consequently, with the onset of Fall rain events, spawner surveys are not practical in the lower river even at moderate water levels as visibility is generally reduced to less than half a metre.

The escapement levels for pink, chum, coho and chinook have shown an increase in recent years over long-term averages but there are several points that must be considered in the current situation.

First, the methodology of estimating escapement on the Englishman changed in 1999. Prior to this year the escapements were estimated by DFO charter patrol and Fisheries Officers and, considering the other demands on the time of these workers, we may assume estimates were based on fewer and less extensive counts. Assessment effort on the Englishman River significantly increased in 1999. When the historic data is examined this change in survey effort must be kept in mind. Any inferences about population trends may be the result of changes in methodology and not necessarily real.

Second, there have been major shifts and increases in commercial and sport fishing restrictions that have a direct influence in the number of salmon returning to fresh water to spawn. In 1998 the troll fishery along the west coast of Vancouver Island was halted, resulting in a dramatic increase in coho escapement along both sides of Vancouver Island in 1998. The progeny of this brood returned in 2001 and continued the large escapement record for this brood line. Additionally, the sport fishing sector has had severe restrictions in both coho and chinook retention, resulting in additional escapement (Baillie et al. 1999, Simpson et al. 1999).

Finally, ocean survivals of smolts, particularly coho and steelhead, have been low in recent years (DFO 2002). Any change in this factor will affect the number of salmon that return to freshwater. This may mask any changes in the population levels of the Englishman stocks that are due to changes in the fresh water habitat. Escapement enumerations are a necessary but not sufficient evaluation of the status of Englishman River stocks and the effectiveness of the recovery plan.

### ***Percent Coverage***

Distribution of salmon varies by species throughout the watershed as the spawning season progresses based on differing run timing and preferred habitat. For example, the majority of any coho that have started to entered freshwater in September would likely

still be holding in the lower river at that time, therefore minimal expansion should be applied to coho counted in the first two surveys. Therefore the peak count of 281 from the 17<sup>th</sup> October survey, covering only the lower reach, was estimated to comprise 95% of the coho in the river at that time. However, by late November most coho will have moved to the upper watershed and tributaries to spawn. Consequently, the final two surveys (2<sup>nd</sup> November and 13<sup>th</sup> January) encompassing only the middle and upper reaches were estimated to have a 75% coverage for coho.

Conversely pink and chum rarely migrate much beyond preferred spawning locations in the lower mainstem, therefore an expansion factor would only need to be applied if this reach was not fully surveyed. This is reflected in the estimated 40% coverage for chum counts on 2<sup>nd</sup> November and 13<sup>th</sup> January when the lower reach was not surveyed.

### ***Pink***

Considering that 1.3 million fry were released in 2004 the pink return to the Englishman river was exceptionally poor in 2006. Although the first survey on 8<sup>th</sup> September was likely passed the normal peak return for pinks there were no carcasses present to indicate an earlier run that may have been missed by this initial survey. The relationship between juvenile production and subsequent adult returns of pinks is not well understood and it is not unusual to see this kind of poor escapement from a seemingly healthy brood year. Pink salmon frequently display this inconsistency between juvenile production and adult returns.

The Englishman pink salmon stock has been supplemented since 1993 by the Quinsam River Hatchery (Bocking and Gaboury, 2001). In subsequent years fry releases have been 1.5 million in 2001, 0.9 million in 2002, 1.5 million in 2003, 1.3 million for 2004 and 0.9 million for 2005. Although between 0.5 million and 1.5 million pink fry have been released each year, it was only the last few years that resulted in any substantial adult return. The resulting progeny from these recent stronger returns should assist in establishing a pink run.

### ***Chinook***

Chinook salmon production has been supplemented by the Englishman River Enhancement Society for nine years. Chinook fry transferred from the Little Qualicum hatchery are reared to the three month stage and released into a pond that drains into the Timberwest side channel. The chinook fry migrate downstream immediately although some individuals reside for one year before smolting. Usually 150,000 to 300,000 fry are released although in 2000 680,000 fry were released.

None of the releases prior to 2003 were marked so the escapement cannot be divided into wild and enhanced origin. Starting in 2003 a thermal mark has been applied so that enhanced chinook will be distinguishable from the wild production when this brood starts to return in 2006.

### ***Chum***

There has been no enhancement of chum in the Englishman River therefore the escapement estimate for chum is indicative of natural production for this system.

Calculating total chum escapement from the 2006 survey data was complex as only two of the five surveys in which chum were counted covered the lower reach. This was one of the primary areas in which chum were observed in 2001 and 2002 (pers. comm., S.



Baillie, Fisheries and Oceans Canada, Nanaimo). As mentioned above, surveys of the lower reach were not always possible due to reduced visibility after rain events or periods of higher run-off.

### **Coho**

Coho production has been supplemented in 1998 and 2001 when 6,000 fed fry and 50,400 fed fry were released into Morison Creek. There has been no enhancement since 2001.

The peak count of 281 coho occurred relatively early in the Fall of 2006, however this may be more a reflection of the extended period starting in early November when no surveys were possible. Coho tend to move into coastal streams during storm events (pers. comm., S. Baillie, Fisheries and Oceans Canada, Nanaimo) and due to the severity and frequency of the Fall storms in 2006 no further surveys were possible once they had commenced. Consequently a portion of the coho escapement may well have been missed during this period of high and turbid river flows. However, low returns of coho were also observed in other Georgia Strait systems in 2006 (anecdotal reports).

## Recommendations

1. An assessment of the success of habitat improvements funded by the PSEF should be conducted. This will require smolt and adult enumeration estimates of coho and steelhead production in the Englishman River.
2. The escapement surveys for 2006 were successful, but they would benefit by having additional swims conducted towards the end of the salmon migration period, and with stream walks added for the tributaries.
3. A survey life study should be conducted as this would improve accuracy of the AUC estimates.
4. Thermally marked chinook will be returning in 2007, therefore it would be worthwhile conducting a carcass-pitch program to recover samples. This would provide an estimate of enhancement contribution to the Englishman River chinook production.

## References cited

- Baillie, S. J., Patten, B., Till, J., Simpson, K., Luedke, W., and Tschaplinski, P. 1999. Assessment of coho stocks on the west coast of Vancouver Island, 1998. Can. Stock Assess. Secret., Res. Doc. 99/166. 43 p.
- Baillie, S. J., and Young, C., 2002. Salmon Escapement to Englishman River, 2001. Fisheries and Oceans Canada unpublished report. 23 pp.
- Baillie, S. J., and Young, C. 2003. Salmon Escapement to Englishman River, 2002. Fisheries and Oceans Canada unpublished report. 21 pp.
- Bocking, R. C., and Gaboury, M. 2001. Englishman River Watershed Recovery Plan, prepared for Pacific Salmon Endowment Fund Society. LGL Ltd. 9768 Second St, Sidney, BC. V8L 3Y8. v + 46 pp.
- DFO, 2002. Coho Salmon in Georgia Basin. DFO Science Stock Status Report D6-07 (2002).
- English, K. K., Bocking, R. C., and Irvine, J. R. 1992. A robust procedure for estimating salmon escapement based on the area-under-the-curve method. Can. J. Fish. Aquat. Sci. 49: 1982-1989.
- Simpson, K., Semple, R., Baillie, S. J., Adkins, B. and Lehmann, S. 1999. Status of coho salmon stocks adjacent to the Strait of Georgia, including the lower Fraser River. Can. Stock Assess. Secret., Res. Doc. 99/133. 74 p.

# Figures




Figure 1. The Englishman River watershed and its relative location on Vancouver Island. Map obtained from the British Columbia Conservation Foundation web site (<http://www.bccf.com/steelhead/focus7.htm#eng> May, 2006).




Figure 2. Map of lower Englishman River showing the three survey reaches used in 2005 escapement surveys.




Figure 3. Water levels for Englishman River from September 1<sup>st</sup>, 2006 to January 31<sup>st</sup>, 2007.

## Tables

Table 1. Salmon escapement estimates to the Englishman River, 1953-2005, from DFO NuSEDS database. UNK = unknown, NO = none observed, NI = not inspected, AP = adults present.

	<b>Sockeye</b>	<b>Coho</b>	<b>Pink</b>	<b>Chum</b>	<b>Chinook</b>
<b>2005</b>	3	3700	4900	950	7300
<b>2004</b>	NI	NI	NI	NI	NI
<b>2003</b>	NO	3200	AP	34800	260
<b>2002</b>	4	3100	12100	9500	600
<b>2001</b>	11	8000	13500	10400	2900
<b>2000</b>	25	5280	1600	3500	1200
<b>1999</b>	20	2978	2500	25000	750
<b>1998</b>	UNK	1500	350	8000	UNK
<b>1997</b>	UNK	200	100	8000	20
<b>1996</b>	UNK	250	800	900	50
<b>1995</b>	UNK	UNK	UNK	2000	UNK
<b>1994</b>	NO	1150	NO	5500	NO
<b>1993</b>	30	246	UNK	1100	24
<b>1992</b>	UNK	440	2000	3500	40
<b>1991</b>	15	800	50	250	50
<b>1990</b>	10	1050	UNK	800	100
<b>1989</b>	30	200	UNK	1500	UNK
<b>1988</b>	30	250	UNK	3000	NO
<b>1987</b>	50	200	UNK	600	NO
<b>1986</b>	10	65	NO	2000	NO
<b>1985</b>	UNK	UNK	UNK	2500	UNK
<b>1984</b>	UNK	2000	UNK	2500	UNK
<b>1983</b>	UNK	UNK	UNK	200	UNK
<b>1982</b>	18	1000	3	2500	14
<b>1981</b>	UNK	300	UNK	400	NO
<b>1980</b>	UNK	300	100	1000	UNK
<b>1979</b>	UNK	1200	UNK	4000	UNK
<b>1978</b>	300	1500	10	6000	75
<b>1977</b>	25	1500	25	1500	25
<b>1976</b>	25	750	25	1500	25
<b>1975</b>	25	400	75	750	75
<b>1974</b>	25	1500	25	5000	25
<b>1973</b>	75	750	25	7500	75
<b>1972</b>	25	400	25	15000	75
<b>1971</b>	25	1500	25	3500	75
<b>1970</b>	25	1500	75	3500	75
<b>1969</b>	25	400	25	7500	75
<b>1968</b>	75	1000	100	6000	115
<b>1967</b>	20	285	NO	500	75
<b>1966</b>	25	1500	200	7500	25
<b>1965</b>	UNK	1500	NO	1500	75
<b>1964</b>	25	1500	NO	1500	25
<b>1963</b>	UNK	750	2	750	25
<b>1962</b>	NO	750	NO	3500	UNK
<b>1961</b>	25	750	25	3500	25
<b>1960</b>	25	400	200	3500	25
<b>1959</b>	1	750	1	3500	UNK
<b>1958</b>	25	750	400	15000	UNK
<b>1957</b>	25	3500	3500	7500	UNK
<b>1956</b>	25	1500	400	750	UNK
<b>1955</b>	25	750	750	1500	UNK
<b>1954</b>	UNK	1500	750	15000	UNK
<b>1953</b>	UNK	750	200	15000	UNK

Table 2. Salmon counts for the three sections of the Englishman River mainstem in 2006. NS = Not Surveyed.

**Pink**

Date	Upper	Middle	Lower	Total
8-Sep-06	NS	NS	47	47
29-Sep-06	NS	1	6	7
17-Oct-06	NS	0	0	0
2-Nov-06	0	0	NS	0
13-Jan-07	NS	0	NS	0

**Chinook**

8-Sep-06	NS	NS	0	0
29-Sep-06	NS	4	145	149
17-Oct-06	NS	66	131	197
2-Nov-06	2	9	NS	11
13-Jan-07	0	0	NS	0

**Chum**

8-Sep-06	NS	NS	0	0
29-Sep-06	NS	0	5	5
17-Oct-06	NS	4	40	44
2-Nov-06	0	2	NS	2
13-Jan-07	0	0	NS	0

**Coho**

8-Sep-06	NS	NS	0	0
29-Sep-06	NS	0	56	56
17-Oct-06	NS	0	281	281
2-Nov-06	3	3	NS	6
13-Jan-07	0	0	NS	0



Table 3. Salmon counts with Observer Efficiency and Percent Coverage expansions for 2006 swim surveys.

**Pink**

Date	Counts	Observer Efficiency	Coverage	System Estimate
8-Sep-06	47	90%	100%	52
29-Sep-06	7	80%	100%	9
17-Oct-06	0	80%	100%	0
2-Nov-06	0	80%	10%	0
13-Jan-07	0	100%	10%	0

**Chinook**

Date	Counts	Observer Efficiency	Coverage	System Estimate
8-Sep-06	0	100%	100%	0
29-Sep-06	172	75%	100%	229
17-Oct-06	197	75%	95%	276
2-Nov-06	12	70%	50%	34
13-Jan-07	0	70%	50%	0

**Coho**

Date	Counts	Observer Efficiency	Coverage	System Estimate
8-Sep-06	0	100%	100%	0
29-Sep-06	56	70%	95%	84
17-Oct-06	281	70%	95%	423
2-Nov-06	6	60%	40%	25
13-Jan-07	0	60%	40%	0

**Chum**

Date	Counts	Observer Efficiency	Coverage	System Estimate
8-Sep-06	0	100%	100%	0
29-Sep-06	5	85%	100%	6
17-Oct-06	44	85%	100%	52
2-Nov-06	2	75%	40%	7
13-Jan-07	0	75%	40%	0